

Representing Even Perfect and Near-Perfect Numbers as Sums of Cubes

PADMA BHUSHAN BORAH, PANKAJ JYOTI MAHANTA, AND MANJIL P. SAIKIA

ABSTRACT. Motivated by recent results of Farhi and Ulas we show that for each $n \equiv \pm 1 \pmod{6}$, the Diophantine equation $2^{n-1}(2^n - 1) = x^3 + y^3 + z^3$ has at least three solutions. We also prove results about the representation of even near-perfect numbers with two distinct prime factors as sums of integral cubes.

1. INTRODUCTION

One of the most frequently studied class of numbers in number theory are the *perfect numbers*. A number n is called a perfect number if it is equal to the sum of its proper divisors. More succinctly, if $\sigma(n) = 2n$ then n is called a perfect number, where $\sigma(n)$ denotes the sum of the distinct positive divisors of n . Even perfect numbers have been known since antiquity and they can be characterized completely by the work of Euclid and Euler (see for instance the book of Hardy and Wright [2]). In particular, any even perfect number n can be written as $n := P_p = 2^{p-1}(2^p - 1)$, where p and $2^p - 1$ are both primes. However, to date no odd perfect number has been found. In fact, it is not known whether they exist at all or not.

Motivated by this search for odd perfect numbers, various generalizations of perfect numbers have been studied extensively in the literature (for instance, see the joint work of the third author with Laugier and Sarmah [3] and the references therein). One of these generalizations is the concept of a *near-perfect number*. For a proper divisor d of n , we call n a near-perfect number with *redundant divisor* d if $\sigma(n) = 2n + d$. Pollack and Shevelev [5] and, Ren and Chen [6] found all near-perfect numbers with two distinct prime factors, which we list below.

Type 1. $n = 2^{t-1}(2^t - 2^k - 1)$, where $2^t - 2^k - 1$ is prime, and 2^k is the redundant divisor.

Type 2. $n = 2^p P_p = 2^{2p-1}(2^p - 1)$, where both p and $2^p - 1$ are prime numbers, and $2P_p$ is the redundant divisor.

Type 3. $n = (2^p - 1)P_p = 2^{p-1}(2^p - 1)^2$, where both p and $2^p - 1$ are prime numbers, and $2^p - 1$ is the redundant divisor.

Type 4. This type consists of only the number 40.

Tang, Ren and Li [8] and Tang, Ma and Feng [7] proved that no odd near-perfect numbers exist with three and four distinct odd prime factors respectively. It is an open problem to show that there exist only finitely many odd near-perfect numbers, as well as to classify all the even near-perfect numbers.

Another aspect of modern number theory which has gained a lot of attention is the question of representing integers by sums of powers. In this connection, Waring's problem asks whether each natural number n can be written as a sum of at most s integral powers of k . For instance,

2010 Mathematics Subject Classification. 11A25, 11D25.

Date: January 27, 2020.

Key words and phrases. perfect numbers, near-perfect numbers, Diophantine equations.

The third author is partially supported by the Leverhulme Trust Research Project Grant RPG-2019-083.

it is known that any natural number can be written as the sum of at most 9 integral cubes. (See the excellent book of Nathanson [4].) Recently, Farhi [1] showed that for even perfect numbers this bound can be reduced; he proved that any even perfect number can be written as a sum of at most five positive integral cubes and further conjectured that they can be written as a sum of at most three positive integral cubes. This conjecture was later studied by Ulas [9], who although could not prove it but found that even perfect numbers can be written as the sum of three integral cubes. In this paper, we investigate similar questions for even near-perfect numbers with at most two distinct prime factors as well as improve upon a recent result of Ulas [9, Theorem 2.1]. An excellent, but brief review of the existing literature related to this problem can be found in the introduction of Ulas' paper [9].

Our main results are the following.

Theorem 1.1. *If $n \equiv \pm 1 \pmod{6}$, then the Diophantine equation*

$$P_n = x^3 + y^3 + z^3, \quad (1.1)$$

has at least three solutions in integers x, y and z .

Theorem 1.2. *If n is an even near-perfect number with two distinct prime factors. Then*

- (1) *n can be written as a sum of at most seven integral cubes if n is of Type 1, provided $t \not\equiv 0 \pmod{3}$.*
- (2) *n can be written as a sum of at most five integral cubes if n is of Type 2.*
- (3) *n can be written as a sum of at most six integral cubes if n is of Type 3.*
- (4) *n can be written as a sum of at most four integral cubes if n is of Type 4.*

The rest of the paper is arranged as follows: Section 2 gives the proofs of Theorems 1.1 and 1.2 and we end this note with some remarks in Section 3.

2. PROOF OF THE RESULTS

2.1. Proof of Theorem 1.1. Before we prove Theorem 1.1 we point out that Ulas [9, Theorem 2.1] has already found two solutions for equation (1.1) when $n \equiv \pm 1 \pmod{6}$, which we list below:

$$\begin{aligned} P_{6n+1} &= (2^{n-2}(2^{3n+2} - 21))^3 + (2^{n-2}(2^{3n+2} + 21))^3 - (11 \cdot 2^{2n-1})^3 \\ &= (2^{4n})^3 + (2^{4n})^3 - (2^{2n})^3, \end{aligned} \quad (2.1)$$

$$\begin{aligned} P_{6n+5} &= (2^n(2^{3(n+1)} + 2^{2(n+1)} + 1))^3 + (2^n(2^{3(n+1)} - 2^{2(n+1)} - 1))^3 - (2^{2(n+1)}(2^{2n+1} + 1))^3 \\ &= (2^{2n+1}(2^{2(n+1)} - 2^{n+1} - 1))^3 + (2^{2n+1}(2^{2(n+1)} + 2^{n+1} - 1))^3 - (2^{4n+3})^3. \end{aligned} \quad (2.2)$$

Proof of Theorem 1.1. It is sufficient to give a solution of equation (1.1) for $n \equiv \pm 1 \pmod{6}$ which is different from equations (2.1) and (2.2). This is given below:

$$\begin{aligned} P_{6n+1} &= (2^{2n}(2^{2n} + 2^n - 1))^3 + (2^{2n}(2^{2n} - 2^n - 1))^3 + (2^{2n})^3, \\ P_{6n+5} &= (2^{4n+3})^3 - (2^{2n+1})^3 - (2^{2n+1})^3. \end{aligned}$$

□

Corollary 2.1. *Every even perfect number $n > 28$ can be represented as a sum of three integral cubes in at least three different ways.*

2.2. Proof of Theorem 1.2. If $N = 2^{p-1}(2^p - 1)$ is a perfect number then Farhi [1] showed that for $a = n^2 + n - 1$ and $b = n^2 - n - 1$, if $p = 6k + 1$ for some integer k , then

$$N = (2^{3k}(2^{2k} + 2^k - 1))^3 + (2^{2k}(2^{2k} - 2^k - 1))^3 + (2^{2k})^3, \quad (2.3)$$

and if $p = 6k + 5$ for some integer k , then

$$N = (3 \cdot 2^{4k+1})^3 + (3 \cdot 2^{4k+1})^3 + (2 \cdot 2^{4k+1})^3 + (2^{2k+1}(2^{2k} + 2^k - 1))^3 + (2^{2k+1}(2^{2k} - 2^k - 1))^3. \quad (2.4)$$

To prove the above representations, he used the following elementary identity, which we use freely in the remainder:

$$2q^6 - 2 = (q^2 + q - 1)^3 + (q^2 - q - 1)^3. \quad (2.5)$$

We prove Theorem 1.2 in a series of Lemmas below.

Lemma 2.2. *Every even near-perfect number with two distinct prime factors of Type 1 can be written as the sum of at most seven integral cubes, provided $t \not\equiv 0 \pmod{3}$.*

Proof. We notice that $t > k$, otherwise n is negative. If $t = 2$, then the only possibility is $k = 1$ for which $2^t - 2^k - 1$ is not a prime. If $t = 3$, then n is either 20 or 12, both of which can be written as the sum of five integral cubes¹. If $t = 4$, then n is 104, 88 or 56. Again, all of them can be written as the sum of five integral cubes².

Let $t \geq 5$, and we write n as

$$n = 2^{t-1}(2^t - 1) - 2^{t+k-1}. \quad (2.6)$$

If $t = 6m + 1$ for some integer m , then

$$2^{t-1}(2^t - 1) = (2^{m-2}(2^{3m+2} - 2))^3 + (2^{m-2}(2^{3m+2} + 2))^3 - (11 \cdot 2^{2m-1})^3. \quad (2.7)$$

From equations (2.6) and (2.7) when $t = 6m + 1$, we have

$$n = (2^{m-2}(2^{3m+2} - 2))^3 + (2^{m-2}(2^{3m+2} + 2))^3 - (11 \cdot 2^{2m-1})^3 - 2^{6m+k},$$

which is the sum of at most four, five or seven integral cubes depending on whether $k \equiv 0, 1$ or $-1 \pmod{3}$ respectively.

If $t = 6m - 1$ for some integer m , then

$$2^{t-1}(2^t - 1) = (2^{m-1}(2^{2m} - 2^m - 1))^3 + (2^{m-1}(2^{2m} + 2^m - 1))^3 - (2^{4m-1})^3. \quad (2.8)$$

From equations (2.6) and (2.8) when $t = 6m - 1$, we have

$$n = (2^{m-1}(2^{2m} - 2^m - 1))^3 + (2^{m-1}(2^{2m} + 2^m - 1))^3 - (2^{4m-1})^3 - 2^{6m+k-2},$$

which is the sum of at most five, seven or four integral cubes depending on whether $k \equiv 0, 1$ or $-1 \pmod{3}$ respectively.

If $t = 6m + 2$ for some integer m , then

$$2^{t-1}(2^t - 1) = (2^{4m+1})^3 - (2^{2m})^3 - (2^{2m})^3. \quad (2.9)$$

From equations (2.6) and (2.9) when $t = 6m + 2$, we have

$$n = (2^{4m+1})^3 - (2^{2m})^3 - (2^{2m})^3 - 2^{6m+1+k},$$

which is the sum of at most five, seven or four integral cubes depending on whether $k \equiv 0, 1$ or $-1 \pmod{3}$ respectively.

¹ $20 = 4^3 - 3^3 - 2^3 - 2^3 - 1^3$ and $12 = 2^3 + 1^3 + 1^3 + 1^3 + 1^3$.

² $104 = 5^3 - 3^3 + 2^3 - 1^3 - 1^3$, $88 = 4^3 + 3^3 - 1^3 - 1^3 - 1^3$ and $56 = 7^3 - 6^3 - 4^3 - 2^3 + 1^3$.

If $t = 6m + 4$ for some integer m , then we have

$$n = 2 \cdot (2^{4m+2})^3 - 2^{6m+k+3} - (2^{2m+1})^3,$$

which is the sum of at most four, five or seven integral cubes depending on whether $k \equiv 0, 1$ or $-1 \pmod{3}$ respectively. \square

Lemma 2.3. *Even near-perfect numbers with two distinct prime factors of Type 2 can be written as sum of at most five integral cubes.*

Proof. Let $n = 2^{2p-1}(2^p - 1)$. If $p = 6m - 1$ for some integer m we have

$$n = (2^{4m-2})^3(4 \cdot (2^{2m})^3 - 2^3).$$

The above is readily seen to be the sum of five cubes.

If $p = 6m + 1$ for some integer m , then we have

$$n = 2^{12m}[2 \cdot 2^{6m} + (2 \cdot 2^{6m} - 2)].$$

Now, using equation (2.5) and some simplification we can write the above as

$$n = 2 \cdot (2^{6m})^3 + (2^{6m} + 2^{5m} - 2^{2m})^3 + (2^{6m} - 2^{5m} - 2^{2m})^3.$$

\square

Lemma 2.4. *Even near-perfect numbers with two distinct prime factors of Type 3 can be written as sum of at most six integral cubes.*

Proof. Let $n = 2^{p-1}(2^p - 1)^2 = 2^{p-1}(2^{2p} - 2^{p+1} + 1)$. If $p = 6m + 1$ for some integer m , then we have

$$n = 2^{6m}(2^p(2 \cdot 2^{6m} - 2) + 1).$$

Using equation (2.5) we get

$$n = 2 \cdot 2^{12m}((2^{2m} + 2^m - 1)^3 + (2^{2m} - 2^m - 1)^3) + 2^{6m},$$

which is a sum of five positive integral cubes.

If $p = 6m + 5$, then we have

$$n = 2^{18m+4} - 2^{6m+3}(2 \cdot 2^{6m+6} - 2).$$

Using equation (2.5) we get

$$n = 4 \cdot 2^{18m+12} - 2^{6m+3}((2^{2m+2} + 2^{m+1} - 1)^3 + (2^{2m+2} - 2^{m+1} - 1)^3),$$

which is a sum of six integral cubes. \square

Proof of Theorem 1.2. Part (1) follows from Lemma 2.2. Part (2) follows from Lemma 2.3. Part (3) follows from Lemma 2.4. Part (4) is trivial since $40 = 4^3 - 2^3 - 2^3 - 2^3$. \square

3. REMARKS

There are several natural questions which we can ask (partly motivated by Farhi's conjecture [1, Conjecture 2.2]).

Question 3.1. It is seen that for Types 2 and 3 even near-perfect numbers, when $p = 6m + 1$ for some integer m then n can be written as a sum of five positive integral cubes. Is this also true when $p = 6m + 5$ for some integer m ?

Question 3.2. What can be said about Type 1 even near-perfect numbers when $t = 3m$ for some integer m ?

Question 3.3. What can be said about the representation of other classes of generalized perfect numbers as sums of cubes or higher powers?

REFERENCES

1. Bakir Farhi, *On the representation of an even perfect number as the sum of five cubes*, *Funct. Approx. Comment. Math.* **57** (2017), no. 2, 277–278. MR 3732899
2. G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, sixth ed., Oxford University Press, Oxford, 2008, Revised by D. R. Heath-Brown and J. H. Silverman, With a foreword by Andrew Wiles. MR 2445243
3. Alexandre Laugier, Manjil P. Saikia, and Upam Sarmah, *Some results on generalized multiplicative perfect numbers*, *Ann. Univ. Ferrara Sez. VII Sci. Mat.* **62** (2016), no. 2, 293–312. MR 3570359
4. Melvyn B. Nathanson, *Additive number theory. The classical bases.*, vol. 164, New York, NY: Springer, 1996 (English).
5. Paul Pollack and Vladimir Shevelev, *On perfect and near-perfect numbers*, *J. Number Theory* **132** (2012), no. 12, 3037–3046. MR 2965207
6. Xiao-Zhi Ren and Yong-Gao Chen, *On near-perfect numbers with two distinct prime factors*, *Bull. Aust. Math. Soc.* **88** (2013), no. 3, 520–524. MR 3189304
7. Min Tang, Xiaoyan Ma, and Min Feng, *On near-perfect numbers*, *Colloq. Math.* **144** (2016), no. 2, 157–188. MR 3493818
8. Min Tang, Xiao-Zhi Ren, and Meng Li, *On near-perfect and deficient-perfect numbers*, *Colloq. Math.* **133** (2013), no. 2, 221–226. MR 3145514
9. Maciej Ulas, *A note on the Diophantine equation $2^{n-1}(2^n - 1) = x^3 + y^3 + z^3$* , *Funct. Approx. Comment. Math.* **60** (2019), no. 1, 87–96. MR 3932606

KESHAB NAGAR, K. B. ROAD, NORTH LAKHIMPUR, ASSAM 787001, INDIA
E-mail address: padmabhushanborah@gmail.com

GONIT SORA, DHALPUR, ASSAM 784165, INDIA
E-mail address: pankaj@gonitsora.com

SCHOOL OF MATHEMATICS, CARDIFF UNIVERSITY, CARDIFF CF24 4AG, UK
E-mail address: SaikiaM@cardiff.ac.uk, manjil@gonitsora.com